



Soils Map



Soils data provided by USDA and NRCS.



State: **Arkansas**
 County: **Lee**
 Location: **34° 44' 52.54, -90° 31' 24.17**
 Township: **Council**
 Acres: **520**
 Date: **10/15/2019**



Maps Provided By:



Area Symbol: AR077, Soil Area Version: 18
 Area Symbol: MS143, Soil Area Version: 12

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Non-Irr Class *c | Common bermudagrass | Cor n | Cotton lint | Pasture | Soybeans | Wheat | *n NCCPI Soybeans |
|-------------------------|--|--------|------------------|----------------------|------------------|---------------------|------------|-------------|------------|-------------|-------------|-------------------|
| 271 | Sharkey clay, 0 to 2 percent slopes, frequently flooded | 140.95 | 27.1% | | IVw | 4.5 | | | | 30 | 45 | 1 |
| 250 | Newellton-Tunica complex, undulating, frequently flooded | 115.68 | 22.2% | | Vw | 4.5 | | | | | | 25 |
| Sk | Sharkey soils, 0 to 1 percent slopes, frequently flooded | 71.49 | 13.7% | | IVw | 4.5 | | | | 30 | 45 | 6 |
| Bs | Bruno soils, frequently flooded | 53.22 | 10.2% | | Vw | | | | 4.5 | | | 18 |
| 73 | Bruno loamy fine sand, 2 to 8 percent slopes, frequently flooded | 47.44 | 9.1% | | Vw | | | | 4.5 | | | 18 |
| 91 | Crevasse sand, 2 to 5 percent slopes, frequently flooded | 38.22 | 7.4% | | Vw | | | | 5 | | | 1 |
| 600 | Water | 21.69 | 4.2% | | | | | | | | | 0 |
| BrB | Bruno fine sandy loam, gently undulating | 11.97 | 2.3% | | III s | | 50 | 400 | 5 | | 30 | 31 |
| 272 | Sharkey clay, ponded | 8.21 | 1.6% | | VII w | | | | | | | 1 |
| W | Water | 6.74 | 1.3% | | | | | | | | | 0 |
| Nf | Newellton soils, frequently flooded | 4.39 | 0.8% | | IVw | 5.5 | | | | 32 | | 26 |
| Weighted Average | | | | | | 2.9 | 1.2 | 9.2 | 1.4 | 12.5 | 19.1 | *n 11.2 |

*n: The aggregation method is "Weighted Average using major components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.