



| | | | | | | | | | | | | | | | | |
|-------------------------|---|-------|------|--|------|------------|---|-------------|------------|--------------|--------------|--|------------|------------|------------|-------------|
| 9 | Ochlockonee sandy loam, occasionally flooded | 54.21 | 2.4% | | Ilw | | | 110 | 22 | | | | 8 | 75 | | 40 |
| 3C3 | Lexington silt loam, 5 to 8 percent slopes, severely eroded | 30.93 | 1.4% | | IVe | | | 60 | | 400 | | | | | 5.5 | 25 |
| 2C | Providence silt loam, 5 to 8 percent slopes, moderately eroded | 30.50 | 1.3% | | IIIe | 8.5 | | 90 | | 750 | | | 9.5 | | | 40 |
| 16 | Cascilla silt loam, 0 to 2 percent slopes, occasionally flooded | 13.27 | 0.6% | | Ilw | 7.5 | 8 | 110 | | 850 | | | 11 | | | 40 |
| 4B | Loring silt loam, 2 to 5 percent slopes, moderately eroded, central | 11.70 | 0.5% | | IIIe | | | 95 | | 750 | | | | | 8 | 40 |
| W | Water | 10.84 | 0.5% | | | | | | | | | | | | | |
| 4C | Loring silt loam, 5 to 8 percent slopes, moderately eroded, central | 3.99 | 0.2% | | IVe | | | 85 | | 700 | | | | | 7.5 | 35 |
| Weighted Average | | | | | | 1.4 | | 28.6 | 0.5 | 168.6 | 596.4 | | 2.5 | 1.8 | 2.7 | 13.6 |

*n: The aggregation method is "Weighted Average using major components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.